

Transcranial Cerebellar Direct Current Stimulation Enhances Verb Generation but Not Verb Naming in Poststroke Aphasia

Paola Marangolo^{1,2}, Valentina Fiori², Carlo Caltagirone^{2,3},
Francesca Pisano¹, and Alberto Priori⁴

Abstract

■ Although the role of the cerebellum in motor function is well recognized, its involvement in the lexical domain remains to be further elucidated. Indeed, it has not yet been clarified whether the cerebellum is a language structure per se or whether it contributes to language processing when other cognitive components (e.g., cognitive effort, working memory) are required by the language task. Neuromodulation studies on healthy participants have suggested that cerebellar transcranial direct current stimulation (tDCS) is a valuable tool to modulate cognitive functions. However, so far, only a single case study has investigated whether cerebellar stimulation enhances language recovery in aphasic individuals. In a randomized, crossover, double-blind design, we explored the effect of cerebellar tDCS coupled with lan-

guage treatment for verb improvement in 12 aphasic individuals. Each participant received cerebellar tDCS (20 min, 2 mA) in four experimental conditions: (1) right cathodal and (2) sham stimulation during a verb generation task and (3) right cathodal and (4) sham stimulation during a verb naming task. Each experimental condition was run in five consecutive daily sessions over 4 weeks. At the end of treatment, a significant improvement was found after cathodal stimulation only in the verb generation task. No significant differences were present for verb naming among the two conditions. We hypothesize that cerebellar tDCS is a viable tool for recovery from aphasia but only when the language task, such as verb generation, also demands the activation of nonlinguistic strategies. ■

INTRODUCTION

During the past two decades, converging neuroscientific evidence has largely documented that the human cerebellum contributes to a much wider range of higher-level cerebral functions than previously accepted. Indeed, although, traditionally, there has been a unanimous agreement that the cerebellum is primarily involved in autonomic and somatic motor processes (Leiner, 2010; Schmahmann, 2010; Strick, Dum, & Fiez, 2009; De Smet, Baillieux, De Deyn, Mariën, & Paquier, 2007; Holmes, 1939), particularly after aphasia reports, there has been a rapidly increasing interest in the cerebellum's role in cognition (Reeber, Otis, & Sillitoe, 2013; Manto & Haines, 2012; Strick et al., 2009). Indeed, several linguistic disorders after acquired cerebellar lesions have been documented (De Smet et al., 2007), such as impaired verbal fluency (Meinzer, Yetim, McMahon, & de Zubicaray, 2016; Stoodley & Schmahmann, 2009; Richter et al., 2007; Leggio, Silveri, Petrosini, & Molinari, 2000; Schmahmann & Sherman, 1998; Molinari, Leggio, & Silveri, 1997; Appollonio, Grafman, Schwartz, Massaquoi, & Hallett, 1993; Akshoomoff,

Courchesne, Press, & Iragui, 1992), agrammatism (Schmahmann & Sherman, 1998; Molinari et al., 1997; Mariën et al., 1996), and naming difficulties (Fabbro, Moretti, & Bava, 2000; Gasparini et al., 1999; Schmahmann & Sherman, 1998). On the basis of these findings, some authors have assumed that the cerebellum represents an “inter-area functional coordinator subserving precisely timed sequential organization of verbal sentences” (Zettin, Cappa, D’amico, Rago, & Perino, 1997; Silveri, Leggio, & Molinari, 1994). Indeed, this process might be compromised in patients with cerebellar lesions (Molinari et al., 1997). Several other cases of aphasia, predominantly, as a result of right cerebellar lesions, characterized by prevailing verbal fluency disturbances, have been described (Mariën & Beaton, 2014; Stoodley & Schmahmann, 2009; Mariën, Engelborghs, Pickut, & De Deyn, 2000; Gasparini et al., 1999). The frequent co-occurrence of a right cerebellar lesion and aphasia led some authors to hypothesize the existence of a “lateralized linguistic cerebellum” (Mariën et al., 1996, 2000, 2014). According to Mariën et al. (1996, 2000, 2014), the aphasic disorder reflects a “diaschisis” phenomenon whereby the damage of the right cerebellum causes a hypofunction of the left frontal cortical areas, “home” of our language representation (Mariën et al., 1996, 2000, 2014). The cerebellum would thus have a role

¹Università Federico II, Naples, Italy, ²IRCCS Fondazione Santa Lucia, Rome, Italy, ³Università degli Studi di Roma Tor Vergata, Rome, Italy, ⁴Università degli Studi di Milano, Milan, Italy

in linguistic representation but only through its connections with the left frontal cerebral language areas (but see Gasparini et al., 1999). In line with this hypothesis, several neuroimaging reports on healthy participants have confirmed the activation of the posterior lateral area of the right cerebellum together with an activation of the left frontal cortex during different linguistic tasks (Chen, Ho, & Desmond, 2014; Stoodley & Schmahmann, 2009; McDermott, Petersen, Watson, & Ojemann, 2003; Gurd et al., 2002; Ojemann et al., 1998; Schlösser et al., 1998), but many conclusions about the role of the cerebellum in language originate from applying word generation tasks (Stoodley, Valera, & Schmahmann, 2010, 2012; Frings et al., 2006; Petersen, Fox, Posner, Mintun, & Raichle, 1989). Petersen and coworkers (1989) reported the first nonmotor linguistic PET activation study in which participants were requested to produce a verb semantically associated to a presented noun. In contrast to the control condition in which the nouns only had to be read or merely repeated, the verb generation condition activated the right lateral cerebellum and a number of left frontal regions. Indeed, this task, which reflects the capacity to generate words according to a given semantic category, requires a large amount of cognitive effort, and it is generally considered to depend on a close cooperation between verbal, executive, and working memory functions, which rely on frontal lobes (Stoodley & Schmahmann, 2009; Bellebaum & Daum, 2007; Gottwald, Mihajlovic, Wilde, & Mehdorn, 2003; Schmahmann & Sherman, 1998; Grafman et al., 1992). Thompson-Schill and colleagues (1998) have suggested that verb generation is a more “difficult” task than naming because it requires the selection of a response from among multiple competitors due to the association strength between an object (noun) and its corresponding verbs (e.g., knife → “cut,” “spread,” “sharp,” “stab”).

Despite variations on the original task design, several other studies have consistently reproduced activation of the right lateral cerebellum during word generation tasks (Grabowski et al., 1996; Martin, Haxby, Lalonde, Wiggs, & Ungerleider, 1995; Raichle et al., 1994). Leiner, Leiner, and Dow (1989) interpreted the simultaneous activation of the right cerebellum and the left Broca’s language area during word generation as “the reflection of accelerated transmission of signals between these two centers during word finding.” Consistent with this assumption, some neuroimaging studies on aphasic individuals have provided evidence in favor of a close connection between the right cerebellar activity and the activation of the contralateral left frontal regions. Indeed, aphasic patients showed an abnormal response from the right cerebellum due to the absence of inputs from the damaged left frontal regions (Connor et al., 2006) and a reactivation of the right cerebellar area after language recovery due to a recruitment of the left perilesional frontal cortex (Marangolo et al., 2016; Heath et al., 2013; Szaflarski, Allendorfer, Banks, Vannest, & Holland, 2013).

Parallel to this increasing interest in the role of the cerebellum in cognition, in more recent years, non-invasive brain stimulation techniques, such as transcranial direct current stimulation (tDCS), have been used to modulate cognitive functions, such as working memory, attention, and language (Lefaucheur et al., 2016; Nitsche & Paulus, 2011).

The assumption on which tDCS is based is that a constant, weak, and continuous current is able to alter the firing rate of the neurons. It has been proposed that cerebellar tDCS is most likely to produce its effects by polarizing Purkinje cells (see Van Dun, Bodranghien, Mariën, & Manto, 2016; Pope & Miall, 2014) and changing the levels/pattern of activity in the deep cerebellar output nuclei, thereby also affecting distant plasticity in human cortical areas (Grimaldi et al., 2016; Van Dun et al., 2016). Indeed, it has been shown that, whereas anodal stimulation, through its excitatory effects, increases the discharge from the Purkinje cells, augmenting the inhibition of the facilitatory pathways from the cerebellar nuclei to the cerebral cortex, cathodal stimulation exerts the opposite effect, through a disinhibition of Purkinje cells, and activates the frontal cerebral cortex (Pope, 2015; Pope & Miall, 2012; Galea & Celnik, 2009).

Pope and Miall (2012) have suggested that one crucial factor for cerebellar tDCS impact is task difficulty. In their study, three groups of 22 participants each performed the paced auditory serial addition task and a variant of this task called the paced auditory serial subtraction task (PASST), together with a verb generation task, before and after anodal, cathodal, or sham tDCS over the right cerebellum. The authors reported an effect on the difficult PASST but not on the easier paced auditory serial addition task. Interestingly, an improvement in the PASST and a reduction in verbal response latencies in verb generation were observed after cathodal right cerebellar tDCS, whereas no effect of anodal stimulation was found (Pope & Miall, 2012). According to Pope and Miall (2012), right cerebellum stimulation has influenced working memory and attention abilities differently depending on task difficulty. Thus, the cerebellum is capable of releasing cognitive resources by disinhibition of the left prefrontal regions, enhancing performance only when the task is cognitively demanding (Pope & Miall, 2012).

Contrary to these findings, in a group of healthy participants, Turkeltaub, Swears, D’Mello, and Stoodley (2016) showed that both anodal and cathodal stimulation over the right cerebellum improves word generation but the effects were found using a different task, namely, a phonemic fluency task. Following Pope and Miall’s suggestion (Pope & Miall, 2012, 2014), the authors hypothesized that cerebellar tDCS did not act directly on the language function per se but on the executive control and response selection components required by the generation task (Turkeltaub et al., 2016).

To date, only a single case study has investigated whether cerebellar tDCS leads to recovery from aphasia.

In a patient with a large bilateral frontoparietal and insular infarct, Sebastian et al. (2016) found that both anodal and sham coupled with language treatment resulted in improved spelling to dictation for trained and untrained words immediately after and 2 months post-treatment but the improvement was greater with anodal tDCS than with sham especially for untrained items. Although the results are interesting and suggest a therapeutic potential of cerebellar tDCS for language recovery, we believe that any final conclusion deserves further investigations. Indeed, as the authors also pointed out, a crucial limitation of their study was that it includes a single case with a large bilateral damage, which is not a lesion typically observed in the aphasic population. In addition, in their experimental design, active tDCS followed sham; thus, any extra benefits of tDCS might be due to having a second treatment after already having had the first treatment (Sebastian et al., 2016).

In this study, we aimed to verify the role of cerebellar tDCS in language processing in a group of 12 aphasic participants with left unilateral damage by contrasting two different language tasks with different demands in terms of cognitive effort: a verb naming (VN) task and a verb generation task. Indeed, with respect to VN in which the production of the correct answer is facilitated by the presented picture, verb generation, because of some combination of both retrieval and competition demands (Snyder, Banich, & Munakata, 2011), relies on different cognitive strategies (Ackermann, Mathiak, & Riecker, 2007; Justus, Ravizza, Fiez, & Ivry, 2005).

Because cathodal stimulation, reducing the inhibition of the Purkinje cells, favors increased excitability of the left frontal language areas (Pope & Miall, 2014; Connor

et al., 2006), in the present work, two experimental conditions were used: right cathodal and sham cerebellar tDCS. On the basis of previous findings, we expected to find that cathodal stimulation would lead to a greater improvement in verb retrieval with respect to the sham condition only in the verb generation task.

METHODS


Participants

Twelve left-brain-damaged participants (six men and six women) with chronic aphasia were included in the study (see Figure 1). Inclusion criteria were native Italian speaker, premorbid right-handedness (Oldfield, 1971), a single left-hemispheric stroke at least 6 months before the investigation, mild nonfluent aphasia with no articulatory difficulties, preserved basic comprehension skills (so as to allow them to be engaged in verbal exchanges with the therapist), and no attentive or memory deficits that might bias their performance. The data analyzed in the current study were collected in accordance with the Declaration of Helsinki and the institutional review board of the IRCCS Fondazione Santa Lucia, Rome, Italy. Before participation, all patients signed informed consent forms.

Clinical Data

The aphasic disorders were assessed using standardized language tests (the Battery for the Analysis of Aphasic Disorders test; Miceli, Laudanna, Burani, & Capasso, 1994; De Renzi & Vignolo, 1962). All patients were classified as nonfluent aphasics as they had reduced verbal output

Figure 1. Brain parenchyma overlap across patients. Color bar refers to the amount of saved voxels, implying 0% being related to the total absence of tissue and 100% being related to the total presence of tissue. As shown, lesion extent included the temporal lobe, the inferior frontal gyrus, the insula, and, partially, the postcentral and precentral gyrus. Axial coordinates refer to the standard space (MNI152).



in spontaneous speech. Their utterances were short, and they were mainly characterized by omissions of verbs as well as errors in verb inflection. Their basic comprehension skills were preserved, but they still have difficulty in comprehending complex materials (mean = 17/36, cutoff score = 29/36, <29 impaired performance on the Token test; De Renzi & Vignolo, 1962). In the noun and VN task, moderate-to-severe word finding difficulties were still present. A neuropsychological battery of tests was also administered, which excluded the presence of attention (i.e., alertness, sustained and selective attention), working memory (i.e., digit span), and executive function deficits that might confound the data (Zimmermann & Fimm, 1994; Spinnler & Tognoni, 1987; see Table 1).

Materials

Sixty pictures of verbs (i.e., to sing, to write) for the VN task and 60 nouns associated to a correspondent verb (i.e., pen → to write) for the verb generation task were selected. For each task, the 60 stimuli were subdivided into two lists of 30 items, each matched for frequency (VN list 1: mean = 28, $SD = 31$; VN list 2: mean = 28, $SD = 31$; nouns list 1: mean = 33, $SD = 32$; nouns list 2: mean = 32, $SD = 33$; unpaired t test, $p > .05$ for each comparison) and length (VN list 1: mean = 8, $SD = 1$; VN list 2: mean = 8, $SD = 2$; nouns list 1: mean = 7, $SD = 2$; nouns list 2: mean = 7, $SD = 2$; unpaired t test, $p > .05$ for each comparison; Bertinetto et al., 2005). The lists were also matched for imageability (estimated on the basis of a sample of 30 healthy participants along a 7-point scale, from 1 = *no imageability* to 7 = *clear imageability* [VN list 1: mean = 6, $SD = 1$; VN list 2: mean = 6, $SD = 1$; nouns list 1: mean = 6, $SD = 1$; nouns list 2: mean = 6, $SD = 1$; unpaired t test, $p > .05$ for each comparison]).

The correlations between the above variables were not significant among the lists suggesting that each measure represented an independent attribute (frequency vs. length, VN list 1: $r = -.22$, $p = .25$; frequency vs. imageability, VN list 1: $r = -.13$, $p = .50$; imageability vs. length, VN list 1: $r = -.08$, $p = .66$; frequency vs. length, VN list 2: $r = -.29$, $p = .12$; frequency vs. imageability, VN list 2: $r = -.23$, $p = .23$; imageability vs. length, VN list 2: $r = .03$, $p = .86$; frequency vs. length, nouns list 1: $r = -.28$, $p = .14$; frequency vs. imageability, nouns list 1: $r = -.23$, $p = .23$; imageability vs. length, nouns list 1: $r = .29$, $p = .12$; frequency vs. length, nouns list 2: $r = -.32$, $p = .09$; frequency vs. imageability, nouns list 2: $r = .13$, $p = .50$; imageability vs. length, nouns list 2: $r = .18$; $p = .34$).

Procedure

Cerebellar tDCS

tDCS was applied using a battery-driven EMS (Bologna, Italy) programmable direct current stimulator with a pair of surface-soaked sponge electrodes (5×7 cm). A constant current of 2-mA intensity was applied through the cathode on the right cerebellar cortex, 1 cm under and

4 cm lateral to theinion (approximately comparable with the projection of the cerebellar lobule VII into the scalp) for 20 min, whereas the reference electrode was positioned over the right shoulder on the deltoid muscle (Pope & Miall, 2012). If applied according to safety guidelines, tDCS is considered to be a safe brain stimulation technique with minor adverse effects (Lefaucheur et al., 2016; Fregni et al., 2015). For each task (VN vs. verb generation), two different stimulation conditions were carried out: (1) cathodal and (2) sham. Sham stimulation was performed exactly like the cathodal condition, but the stimulator was turned off after 30 sec (Gandiga, Hummel, & Cohen, 2006). Thus, we had four different experimental conditions: (1) right cathodal cerebellar tDCS for VN, (2) sham for VN, (3) right cathodal cerebellar tDCS for verb generation, and (4) sham for verb generation. For each task, the 60 stimuli were subdivided into two lists of 30 items, each matched for frequency, length, and imageability. The assignment of each list of stimuli was randomized across the two conditions (cathodal vs. sham). All patients underwent the four experimental conditions whose order was randomized across participants. To ensure the double-blind procedure, both the experimenter and the patient were blinded regarding the stimulation condition, and the stimulator was turned on/off by another person. At the end of each experimental condition, participants were asked if they were aware of which condition (real or sham) they were in. We inferred that all participants well tolerated the stimulation by interpreting their spontaneous report as well as the results from a questionnaire completed by the participant at the end of each experimental condition (see Fertoni, Rosini, Cotelli, Rossini, & Miniussi, 2010, for the questionnaire). Itch was the most commonly reported sensation with light (16% of the participants) to moderate (83% of the participants) intensity. Participants reported that the sensation started at the beginning of the stimulation and stopped after a few minutes both during the real and/or sham stimulation. Thus, none of the participants was able to distinguish between the two conditions. A paired t test did not show any significant difference in the participants' perception of sensation between the real and sham conditions ($p > .05$).

Treatment

Once the electrodes were placed, participants performed the two tasks while they received 20 min of cerebellar tDCS. Each stimulation condition was performed in five consecutive daily sessions over 1 week with 6 days of intersession interval. The order of item presentation was randomized across sessions. During the VN task, participants were asked to name aloud each picture that appeared on the PC screen (screen size = 15 in., viewing distance = 1 m) for 20 sec preceded by a fixation point, which lasted 800 msec (see also Fiori et al., 2013, for a similar procedure). If the participant failed or did not

Table 1. Sociodemographic and Clinical Data of the 12 Nonfluent Aphasic Patients

<i>P</i>	Age (Years)	Educ Level (Years)	Time Post Onset	NV (%)	VN (%)	NC (%)	VC (%)	TT (Cutoff = 29/36, <29 Impaired)	Attentional Abilities (Scores in Percentile <5 Impaired)	WM (Cutoff = 5 ± 2, <5 Impaired)	Weigl's Sorting Test (Cutoff = 4,50, <4,50 Impaired)
1	50	16	2 years	48	20	100	100	14/36	Alertness (tot): 60 Sustained att (tot): 75	WM: 4	9,75
2	61	13	1 year 3 months	90	64	100	100	14/36	Selective att (tot): 55 Alertness (tot): 99	WM: 5	10
3	46	8	1 year 10 months	30	20	100	100	14/36	Sustained att (tot): 76 Selective att (tot): 94	WM: 4	10
4	65	13	1 year 7 months	45	40	100	100	15/36	Alertness (tot): 65 Sustained att (tot): 34	WM: 4	9,25
5	68	18	1 year 9 months	77	67	100	100	21/36	Sustained att (tot): 70 Selective att (tot): 60	WM: 5	12,5
6	57	13	2 years 9 months	67	64	100	100	14/36	Alertness (tot): 89 Sustained att (tot): 54	WM: 5	10,75
7	70	13	1 year 9 months	70	65	100	100	21/36	Selective att (tot): 66 Alertness (tot): 35	WM: 5	11,5
									Sustained att (tot): 30 Alertness (tot): 95		
									Sustained att (tot): 80 Selective att (tot): 70		

8	48	18	1 year 2 months	33	40	100	100	14/36	Alertness (tot): 94 Sustained att (tot): 55 Selective att (tot): 50	WM: 5	9,75
9	51	8	1 year 6 months	30	30	100	100	23/36	Alertness (tot): 97 Sustained att (tot): 60 Selective att (tot): 84	WM: 5	11
10	60	13	1 year 6 months	55	60	100	100	23/36	Alertness (tot): 90 Sustained att (tot): 80 Selective att (tot): 60	WM: 6	10
11	61	10	1 year 5 month	43	25	100	100	14/36	Alertness (tot): 78 Sustained att (tot): 81 Selective att (tot): 50	WM: 5	10,5
12	56	13	3 years 1 month	70	71	100	100	14/36	Alertness (tot): 90 Sustained att (tot): 62 Selective att (tot): 85	WM: 6	10,75

P = participants; Educ Level = educational level; NN = noun naming; NC = noun comprehension; VC = verb comprehension (Battery for the Analysis of Aphasic Disorders test; Miceli et al., 1994); TT = Token test, cutoff = 29/36 (De Renzi & Vignolo, 1962); att = attention; WM = working memory. Attentional abilities (Zimmermann & Fimm, 1994): patients' scores are reported in percentile; scores below the fifth percentile are considered impaired (i.e., digit span, Orsini et al., 1987; Spinnler & Tognoni, 1987). Weigl's sorting test (Italian version; Spinnler & Tognoni, 1987): For each patient, raw scores are reported.

answer within 20 sec, the corresponding written name was presented below the picture for 5 sec and the participant was asked to read the word aloud. The pair of stimuli remained on the screen until the participant read the word or 5 sec elapsed. In all cases, participants were able to correctly read the word. During the verb generation task, the examiner orally presented a noun (i.e., trampoline) and participants were required to produce within 20 sec the most appropriate corresponding verb (i.e., to jump). Because previous studies have reported activation of the right lateral cerebellum during a verb generation task independently of the input modality (Richter et al., 2004; Petersen et al., 1989), we chose to orally present the noun to prevent participants from “task errors” such as reading the noun by presenting a written noun (Thompson-Schill et al., 1998) or naming the object by presenting the object picture (Kurland, Reber, & Stokes, 2014).

For both tasks, the examiner manually recorded the response on a separate sheet. If the participant did not respond within the 20-sec interval, the program automatically presented the subsequent picture or noun. Vocal RTs were calculated from the presentation of the picture or the noun to the pronunciation of the first phoneme through Audacity 2.1.2 Software. RTs were recorded only if the participants responded within 20 sec.

Data Analysis

Before the experiment, the two lists of stimuli were presented to a group of 30 healthy individuals (15 men and 15 women) matched for age (40–75 years) and educational level (13–17 years) to the aphasic group. Each participant was asked to produce for each presented verb for the VN task and noun for the verb generation task the most appropriate corresponding verb, with no interference from the examiner. Only those verbs that elicited

at least 80% of agreement among participants in the two lists were considered correct responses and therefore used for a response accuracy analysis in the aphasic group.

Data were analyzed with STATISTICA 10 (StatSoft, Inc., Tulsa, OK). Statistical analyses were performed with two separate ANOVAs, respectively, for response accuracies and vocal RTs with three within-participant factors: Task (verb generation vs. VN), Condition (cathodal stimulation vs. sham), and Time (baseline [T0] vs. end of the treatment [T5] vs. follow-up [FU]). If the ANOVA showed significant effects, respective post hoc Bonferroni tests were conducted.

RESULTS

Accuracy

The analysis showed a significant effect of Condition ($F(1, 11) = 13.88, p < .01$) and Time ($F(2, 22) = 77.94, p < .001$). The interaction Task \times Condition \times Time was also significant ($F(2, 22) = 16.21, p < .001$). Indeed, although all experimental conditions led to a significant greater percentage of correct responses at the end of treatment (T5) compared with the baseline (T0; verb generation: difference between T5 and T0, cathodal = 44%, $p < .001$; sham = 15%, $p = .001$; VN: difference between T5 and T0, cathodal = 15%, $p = .001$; sham = 12%, $p = .007$), at the end of treatment, only in the verb generation task, the percentage of correct responses was greater after cathodal stimulation compared with sham (cathodal vs. sham = 28%, $p < .001$), and this difference persisted at FU (cathodal vs. sham = 25%, $p < .001$). No differences between the two stimulation conditions were found for the naming task (cathodal vs. sham: T5 = 4%, $p = 1$; cathodal vs. sham: FU = 2%, $p = 1$; see Figure 2).




Figure 2. Mean percentage of response accuracy for verb generation and VN task at baseline (T0), at the end of the treatment (T5), and at FU for the cathodal and sham conditions, respectively ($*p \leq .001$, $**p < .01$). Error bars represent SEM.

Table 2. Mean Number of Errors in the Verb Generation and VN Task for Cathodal and Sham tDCS, Respectively ($\pm SD$)

Type of Errors	Verb Generation, Cathodal		Verb Generation, Sham		VN, Cathodal		VN, Sham	
	T0	T5	T0	T5	T0	T5	T0	T5
No responses	19 (± 5)	7 (± 3)	18 (± 6)	13 (± 7)	17 (± 6)	13 (± 6)	18 (± 7)	12 (± 6)
Semantic paraphasias	3 (± 2)	2 (± 1)	3 (± 2)	3 (± 2)	2 (± 1)	2 (± 2)	2 (± 2)	3 (± 2)
Unrelated verb responses	2 (± 2)	2 (± 2)	2 (± 2)	3 (± 3)	1 (± 1)	1 (± 2)	1 (± 1)	1 (± 2)

To further investigate if tDCS had a different impact on the participant's response, we classified the errors made by each participant in all experimental conditions. As shown in Table 2, errors were (1) no responses, (2) semantic paraphasias, and (3) unrelated verb responses, but at baseline (T0), for both the VN task and the verb generation task, errors were predominantly "no responses." Thus, we conducted an ANOVA on the number of "no responses" with three within-participant factors: Task (verb generation vs. VN), Condition (cathodal vs. sham), and Time (baseline [T0] vs. end of the treatment [T5]). The analysis revealed a significant interaction of Task \times Condition \times Time ($F(1, 11) = 8.29, p = .01$). Indeed, although all experimental conditions led to a lower number of "no responses" at the end of treatment (T5) compared with the baseline (T0; verb generation: difference between T5 and T0, cathodal = 12, $p < .001$; sham = 5, $p = .04$; VN: difference between T5 and T0, cathodal = 5, $p = .04$; sham = 6, $p = .03$), at the end of treatment (T5), only in the verb generation task, the number of "no responses" was lower after cathodal stimulation compared with sham (cathodal vs. sham = -6, $p = .02$). No differences between the two stimulation conditions were found for the VN task (cathodal vs. sham: T5 = 1, $p = 1$).

Thus, these results resembled those previously found for the accuracy data.

Vocal RTs

The analysis showed a significant effect of Condition ($F(1, 11) = 6.20, p = .03$) and Time ($F(2, 22) = 40.24, p < .001$). The interaction Task \times Condition \times Time was also significant ($F(2, 22) = 11.12, p < .001$). Indeed, although all experimental conditions led to faster vocal RTs at the end of treatment (T5) compared with baseline (T0; verb generation: difference between T5 and T0, cathodal = 4334 msec, $p < .001$; sham = 1644 msec, $p = .003$; VN: difference between T5 and T0, cathodal = 1897 msec, $p = .001$; sham = 1932 msec, $p < .001$), at the end of treatment, only in the verb generation task, vocal RTs were faster after cathodal stimulation compared with sham (cathodal vs. sham = -2350 msec, $p < .001$), and this difference persisted at FU (cathodal vs. sham = -2250 msec, $p < .001$). No differences between the two stimulation conditions were found for the VN task (cathodal vs. sham: T5 = 10 msec, $p = 1$; cathodal vs. sham: FU = 23 msec, $p = 1$; see Figure 3).




Figure 3. Mean vocal RTs for verb generation and VN at baseline (T0), at the end of the treatment (T5), and at FU for the cathodal and sham conditions, respectively ($*p \leq .001, **p < .01$). Error bars represent SEM.

DISCUSSION

The aim of this study was to investigate whether cerebellar tDCS coupled with language training improves verb retrieval in nonfluent chronic aphasic individuals. Our findings showed that cathodal stimulation differently affected verb recovery depending on the language task. Indeed, at the end of treatment, only the verb generation task led to a significant improvement in verb retrieval. Moreover, FU testing showed that these effects lasted over 1 week after the intervention. This specificity argues against an effect simply due to enhanced cognitive arousal, which should have influenced both language tasks.

As stated in the Introduction, several studies have already supported the hypothesis that the cerebellum plays a role in language processing but depends on task demands (Pope & Miall, 2012, 2014; Stoodley et al., 2010, 2012; Ackermann et al., 2007). Ackermann et al. (2007) have argued that nonlinguistic aspects of task performance, such as the amount of effort or the degree of automaticity, might account for cerebellar involvement during verb generation tasks. Similarly, Stoodley and Schmahmann (2009) have claimed that the cerebellum takes part not in the language function per se but only when the task is cognitively demanding and therefore engages other cognitive components, such as working memory and/or executive functions (Stoodley & Schmahmann, 2009). Indeed, apart from motor control and higher-order aspects of speech production, a variety of studies point to a contribution of the cerebellum to executive and memory tasks (Ackermann et al., 2007). Because the paradigm of verb generation involves the production and selection of different verbal responses (Thompson-Schill et al., 1998), prearticulatory rehearsal processes are engaged as well, which rely to working memory processes (Ackermann et al., 2007; Helmuth, Ivry, & Shimizu, 1997). Indeed, our choice to directly compare cerebellar tDCS effects in two verb production tasks was made taking into account the substantial differences between the two tasks.

Like confrontation naming, verb generation is a semantic association task in which the participant has to produce a verb strictly associated to a given noun. Much of the cognitive demand between the two tasks is shared, including semantic and lexical retrieval processes and the planning, execution, and monitoring of speech production (e.g., Levelt, 1989). However, whereas verb generation requires the patient to creatively link a noun to a verb choosing among competing response alternatives (Thompson-Schill et al., 1998), in VN, the correct answer is univocally determined by the presented picture and the task is one of the earliest linguistic skills developmentally mastered and thus is an overlearned task (Herholz et al., 1997). Interestingly, although verb generation is a task more cognitive demanding than VN and persons with aphasia generally experience greatest difficulty with verb generation (Martin & Cheng, 2006; Thompson-Schill

et al., 1998), our aphasic patients benefited only for this task after right cerebellar cathodal stimulation.

Although our data are only behavioural, we might speculate that right cathodal cerebellar stimulation, through a disinhibition of the Purkinje cells, has favored the engagement of the left frontal areas, which, in turn, enhanced the activation of executive and memory components required by the verb generation task (Pope & Miall, 2014; Connor et al., 2006; Mariën, Engelborghs, Fabbro, & De Deyn, 2001; Mariën et al., 1996). Indeed, most of our patients had a partial damage to the left frontal areas (see Figure 1); thus, the hypothesis can be advanced that subregions of the left frontal cortex took part in verb recovery. Accordingly, several studies have already shown that the same facilitatory patterns may be observed in a verbal fluency task after cathodal cerebellar tDCS or anodal stimulation over the frontal cortex (Pope & Miall, 2012; Iyer et al., 2005). Confirming evidence for a functional relationship between the left frontal cortex and the cerebellum comes also from a recent study combining bilateral tDCS and resting state fMRI in a group of left-brain-damaged population (Marangolo et al., 2016). Indeed, in nine chronic aphasic patients, Marangolo et al. (2016) found that bilateral anodic stimulation over the left inferior frontal area and cathodal contralesional stimulation over its right homologue coupled with an intensive language treatment led to functional connectivity changes within the left damaged hemisphere, together with the cerebellum (Marangolo et al., 2016). In agreement with our hypothesis that the cerebellum is functionally connected to the language network, very recently, D'Mello, Turkeltaub, and Stoodley (2017) acquired behavioral and resting state fMRI data, during a sentence completion task, before and after cerebellar tDCS in a group of healthy adults. Relative to sham, anodal tDCS increased activation in the right cerebellum only when the preceding context in the sentence modulated the predictability of the target word (predictive sentences). In the same study (D'Mello et al., 2017), functional connectivity changes were also found in the left language areas, including the left inferior frontal gyrus. Thus, these data showed that cerebellar neuromodulation specifically alters activation patterns during semantic prediction tasks (D'Mello et al., 2017). Similarly, in our work, cerebellar tDCS improved the generation of highly predictable verbs semantically associated to the presented nouns.

It might be finally argued that the effects found were an artifact of linguistic variables, so that, at the end of treatment, the verbs produced in the generation task had higher frequency and/or were shorter (in terms of number of phonemes) than the verbs produced in the naming task. However, statistical analyses performed to control for those factors did not show any significant difference between the correct responses given in the two tasks (verb generation: mean frequency = 32, $SD = 9$; mean length = 8, $SD = 1$; VN: mean frequency = 31, $SD = 8$; mean length = 8, $SD = 0$; t tests, $p = .72$ and $p = .69$, respectively, for frequency and length).

In conclusion, although our data deserve further investigations, they suggest that cerebellar tDCS might be a viable tool for enhancing language recovery in chronic aphasia. Because our results point to potential therapeutic benefits of cerebellar stimulation only for complex language tasks, we believe that these findings have important implications for aphasia. Indeed, they address the possibility that the cerebellum supports cognitive functions that are important for language recovery.

Acknowledgments

We are extremely grateful to Dr. Tommaso Gili for his help in drawing Figure 1.

Reprint requests should be sent to Paola Marangolo, Dipartimento di Studi Umanistici, Università degli Studi di Napoli Federico II, Via Porta di Massa, 1, 80133 Napoli, Italia, or via e-mail: paola.marangolo@gmail.com.

REFERENCES

- Ackermann, H., Mathiak, K., & Riecker, A. (2007). The contribution of the cerebellum to speech production and speech perception: Clinical and functional imaging data. *Cerebellum*, *6*, 202–213.
- Akshoomoff, N. A., Courchesne, E., Press, G. A., & Iragui, V. (1992). Contribution of the cerebellum to neuropsychological functioning: Evidence from a case of cerebellar degenerative disorder. *Neuropsychologia*, *30*, 315–328.
- Appollonio, I. M., Grafman, J., Schwartz, V., Massaquoi, S., & Hallett, M. (1993). Memory in patients with cerebellar degeneration. *Neurology*, *43*, 1536–1544.
- Bellebaum, C., & Daum, I. (2007). Cerebellar involvement in executive control. *Cerebellum*, *6*, 184–192.
- Bertinetto, P. M., Burani, C., Laudanna, A., Marconi, L., Ratti, D., Rolando, C., et al. (2005). *Corpus e Lessico di Frequenza dell'Italiano Scritto (CoLFIS)*. Istituto di Linguistica Computazionale, Unità staccata di Genova. http://linguistica.sns.it/CoLFIS/CoLFIS_home.htm.
- Chen, S.-H. A., Ho, M.-H. R., & Desmond, J. E. (2014). A meta-analysis of cerebellar contributions to higher cognition from PET and fMRI studies. *Human Brain Mapping*, *35*, 593–615.
- Connor, L. T., Braby, T. D., Snyder, A. Z., Lewis, C., Blasi, V., & Corbetta, M. (2006). Cerebellar activity switches hemispheres with cerebral recovery in aphasia. *Neuropsychologia*, *44*, 171–177.
- De Renzi, E., & Vignolo, L. A. (1962). The Token test: A sensitive test to detect receptive disturbances in aphasics. *Brain*, *85*, 665–678.
- De Smet, H. J., Baillieux, H., De Deyn, P. P., Mariën, P., & Paquier, P. (2007). The cerebellum and language: The story so far. *Folia Phoniatrica et Logopaedica*, *59*, 165–170.
- D'Mello, A. M., Turkeltaub, P. E., & Stoodley, C. J. (2017). Cerebellar tDCS modulates neural circuits during semantic prediction: A combined tDCS-fMRI study. *Journal of Neuroscience*, *37*, 1604–1613.
- Fabbro, F., Moretti, R., & Bava, A. (2000). Language impairments in patients with cerebellar lesions. *Journal of Neurolinguistics*, *13*, 173–188.
- Fertonani, A., Rosini, S., Cotelli, M., Rossini, P. M., & Miniussi, C. (2010). Naming facilitation induced by transcranial direct current stimulation. *Behavioural Brain Research*, *208*, 311–318.
- Fiori, V., Cipollari, S., Di Paola, M., Razzano, C., Caltagirone, C., & Marangolo, P. (2013). tDCS stimulation segregates words in the brain: Evidence from aphasia. *Frontiers in Human Neuroscience*, *7*, 269.
- Fregni, F., Nitsche, M. A., Loo, C. K., Brunoni, A. R., Marangolo, P., Leite, J., et al. (2015). Regulatory considerations for the clinical and research use of transcranial direct current stimulation (tDCS): Review and recommendations from an expert panel. *Clinical Research and Regulatory Affairs*, *32*, 22–35.
- Frings, M., Dimitrova, A., Schorn, C. F., Elles, H. G., Hein-Kropp, C., Gizewski, E. R., et al. (2006). Cerebellar involvement in verb generation: An fMRI study. *Neuroscience Letters*, *409*, 19–23.
- Galea, J. M., & Celnik, P. (2009). Brain polarization enhances the formation and retention of motor memories. *Journal of Neurophysiology*, *102*, 294–301.
- Gandiga, P. C., Hummel, F. C., & Cohen, L. G. (2006). Transcranial DC stimulation (tDCS): A tool for double-blind sham-controlled clinical studies in brain stimulation. *Clinical Neurophysiology*, *117*, 845–850.
- Gasparini, M., Di Piero, V., Ciccarelli, O., Cacioppo, M. M., Pantano, P., & Lenzi, G. L. (1999). Linguistic impairment after right cerebellar stroke: A case report. *European Journal of Neurology*, *6*, 353–356.
- Gottwald, B., Mihajlovic, Z., Wilde, B., & Mehdorn, H. M. (2003). Does the cerebellum contribute to specific aspects of attention? *Neuropsychologia*, *41*, 1452–1460.
- Grabowski, T. J., Frank, R. J., Brown, C. K., Damasio, H., Ponto, L. L., Watkins, G. L., et al. (1996). Reliability of PET activation across statistical methods, subject groups, and sample sizes. *Human Brain Mapping*, *4*, 23–46.
- Grafman, J., Litvan, I., Massaquoi, S., Stewart, M., Sirigu, A., & Hallett, M. (1992). Cognitive planning deficit in patients with cerebellar atrophy. *Neurology*, *42*, 1493–1496.
- Grimaldi, G., Argyropoulos, G. P., Bastian, A., Cortes, M., Davis, N. J., Edwards, D. J., et al. (2016). Cerebellar transcranial direct current stimulation (ctDCS): A novel approach to understanding cerebellar function in health and disease. *Neuroscientist*, *22*, 83–97.
- Gurd, J. M., Amunts, K., Weiss, P. H., Zafiris, O., Zilles, K., Marshall, J. C., et al. (2002). Posterior parietal cortex is implicated in continuous switching between verbal fluency tasks: An fMRI study with clinical implications. *Brain*, *125*, 1024–1038.
- Heath, S., McMahon, K. L., Nickels, L., Angwin, A., MacDonald, A. D., van Hees, S., et al. (2013). Facilitation of naming in aphasia with auditory repetition: An investigation of neurocognitive mechanisms. *Neuropsychologia*, *51*, 1534–1548.
- Helmuth, L. L., Ivry, R. B., & Shimizu, N. (1997). Preserved performance by cerebellar patients on tests of word generation, discrimination learning, and attention. *Learning and Memory*, *3*, 456–474.
- Herholz, K., Reulen, H. J., von Stockhausen, H. M., Thiel, A., Ilmberger, J., Kessler, J., et al. (1997). Preoperative activation and intraoperative stimulation of language-related areas in patients with glioma. *Neurosurgery*, *41*, 1253–1260.
- Holmes, G. (1939). The cerebellum of man. *Brain*, *62*, 1–30.
- Iyer, M. B., Mattu, U., Grafman, J., Lomarev, M., Sato, S., & Wassermann, E. M. (2005). Safety and cognitive effect of frontal DC brain polarization in healthy individuals. *Neurology*, *64*, 872–875.
- Justus, T., Ravizza, S. M., Fiez, J. A., & Ivry, R. B. (2005). Reduced phonological similarity effects in patients with damage to the cerebellum. *Brain and Language*, *95*, 304–318.

- Kurland, J., Reber, A., & Stokes, P. (2014). Beyond picture naming: Norms and patient data for a verb-generation task. *American Journal of Speech-Language Pathology, 23*, S259–S270.
- Lefaucheur, J.-P., Antal, A., Ayache, S. S., Benninger, D. H., Brunelin, J., Cogiamanian, F., et al. (2016). Evidence-based guidelines on the therapeutic use of transcranial direct current stimulation (tDCS). *Clinical Neurophysiology, 128*, 56–92.
- Leggio, M. G., Silveri, M. C., Petrosini, L., & Molinari, M. (2000). Phonological grouping is specifically affected in cerebellar patients: A verbal fluency study. *Journal of Neurology, Neurosurgery and Psychiatry, 69*, 102–106.
- Leiner, H. C. (2010). Solving the mystery of the human cerebellum. *Neuropsychology Review, 20*, 229–235.
- Leiner, H. C., Leiner, A. L., & Dow, R. S. (1989). Reappraising the cerebellum: What does the hindbrain contribute to the forebrain? *Behavioral Neuroscience, 103*, 998–1008.
- Levelt, W. J. M. (1989). *Speaking: From intention to articulation*. Cambridge, MA: MIT Press.
- Manto, M., & Haines, D. (2012). Cerebellar research: Two centuries of discoveries. *Cerebellum, 11*, 446–448.
- Marangolo, P., Fiori, V., Sabatini, U., De Pasquale, G., Razzano, C., Caltagirone, C., et al. (2016). Bilateral transcranial direct current stimulation language treatment enhances functional connectivity in the left hemisphere: Preliminary data from aphasia. *Journal of Cognitive Neuroscience, 28*, 724–738.
- Mariën, P., Ackermann, H., Adamaszek, M., Barwood, C. H. S., Beaton, A., Desmond, J., et al. (2014). Consensus paper: Language and the cerebellum: An ongoing enigma. *Cerebellum, 13*, 386–410.
- Mariën, P., & Beaton, A. (2014). The enigmatic linguistic cerebellum: Clinical relevance and unanswered questions on nonmotor speech and language deficits in cerebellar disorders. *Cerebellum & Ataxias, 1*, 12.
- Mariën, P., Engelborghs, S., Fabbro, F., & De Deyn, P. P. (2001). The lateralized linguistic cerebellum: A review and a new hypothesis. *Brain and Language, 79*, 580–600.
- Mariën, P., Engelborghs, S., Pickut, B. A., & De Deyn, P. P. (2000). Aphasia following cerebellar damage: Fact or fallacy? *Journal of Neurolinguistics, 13*, 145–171.
- Mariën, P., Saerens, J., Nanhoe, R., Moens, E., Nagels, G., Pickut, B. A., et al. (1996). Cerebellar induced aphasia: Case report of cerebellar induced prefrontal aphasic language phenomena supported by SPECT findings. *Journal of the Neurological Sciences, 144*, 34–43.
- Martin, A., Haxby, J. V., Lalonde, F. M., Wiggs, C. L., & Ungerleider, L. G. (1995). Discrete cortical regions associated with knowledge of color and knowledge of action. *Science, 270*, 102–105.
- Martin, R. C., & Cheng, Y. (2006). Selection demands versus association strength in the verb generation task. *Psychonomic Bulletin & Review, 13*, 396–401.
- McDermott, K. B., Petersen, S. E., Watson, J. M., & Ojemann, J. G. (2003). A procedure for identifying regions preferentially activated by attention to semantic and phonological relations using functional magnetic resonance imaging. *Neuropsychologia, 41*, 293–303.
- Meinzer, M., Yetim, Ö., McMahon, K., & de Zubicaray, G. (2016). Brain mechanisms of semantic interference in spoken word production: An anodal transcranial direct current stimulation (atDCS) study. *Brain and Language, 157–158*, 72–80.
- Miceli, G., Laudanna, A., Burani, C., & Capasso, R. (1994). *Batteria per l'Analisi del Deficit Afasico. BADA [BADA A battery for the assessment of aphasic disorders]*. Roma, Italia: CEPISAG.
- Molinari, M., Leggio, M. G., & Silveri, M. C. (1997). Verbal fluency and agrammatism. *International Review of Neurobiology, 41*, 325–339.
- Nitsche, M. A., & Paulus, W. (2011). Transcranial direct current stimulation—Update 2011. *Restorative Neurology and Neuroscience, 29*, 463–492.
- Ojemann, J. G., Buckner, R. L., Akbudak, E., Snyder, A. Z., Ollinger, J. M., McKinstry, R. C., et al. (1998). Functional MRI studies of word-stem completion: Reliability across laboratories and comparison to blood flow imaging with PET. *Human Brain Mapping, 6*, 203–215.
- Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia, 9*, 97–113.
- Orsini, A., Grossi, D., Capitani, E., Laiacona, M., Papagno, C., & Vallar, G. (1987). Verbal and spatial immediate memory span: Normative data from 1355 adults and 112 children. *Italian Journal of Neuroscience, 8*, 539–548.
- Petersen, S. E., Fox, P. T., Posner, M. I., Mintun, M., & Raichle, M. E. (1989). Positron emission tomographic studies of the processing of single words. *Journal of Cognitive Neuroscience, 1*, 153–170.
- Pope, P. A. (2015). Modulating cognition using transcranial direct current stimulation of the cerebellum. *Journal of Visualized Experiments, 96*, e52302.
- Pope, P. A., & Miall, R. C. (2012). Task-specific facilitation of cognition by cathodal transcranial direct current stimulation of the cerebellum. *Brain Stimulation, 5*, 84–94.
- Pope, P. A., & Miall, R. C. (2014). Restoring cognitive functions using non-invasive brain stimulation techniques in patients with cerebellar disorders. *Frontiers in Psychiatry, 5*, 1–7.
- Raichle, M. E., Fiez, J. A., Videen, T. O., MacLeod, A. M., Pardo, J. V., Fox, P. T., et al. (1994). Practice-related changes in human brain functional anatomy during nonmotor learning. *Cerebral Cortex, 4*, 8–26.
- Reeber, S. L., Otis, T. S., & Sillitoe, R. V. (2013). New roles for the cerebellum in health and disease. *Frontiers in Systems Neuroscience, 7*, 83.
- Richter, S., Gerwig, M., Aslan, B., Wilhelm, H., Schoch, B., Dimitrova, A., et al. (2007). Cognitive functions in patients with MR-defined chronic focal cerebellar lesions. *Journal of Neurology, 254*, 1193–1203.
- Richter, S., Kaiser, O., Hein-Kropp, C., Dimitrova, A., Gizewski, E., Beck, A., et al. (2004). Preserved verb generation in patients with cerebellar atrophy. *Neuropsychologia, 42*, 1235–1246.
- Schlösser, R., Hutchinson, M., Joseffer, S., Rusinek, H., Saarimaki, A., Stevenson, J., et al. (1998). Functional magnetic resonance imaging of human brain activity in a verbal fluency task. *Journal of Neurology, Neurosurgery and Psychiatry, 64*, 492–498.
- Schmahmann, J. D. (2010). The role of the cerebellum in cognition and emotion: Personal reflections since 1982 on the dysmetria of thought hypothesis, and its historical evolution from theory to therapy. *Neuropsychology Review, 20*, 236–260.
- Schmahmann, J. D., & Sherman, J. C. (1998). The cerebellar cognitive affective syndrome. *Brain, 121*, 561–579.
- Sebastian, R., Saxena, S., Tsapkini, K., Faria, A. V., Long, C., Wright, A., et al. (2016). Cerebellar tDCS: A novel approach to augment language treatment post-stroke. *Frontiers in Human Neuroscience, 10*, 695.
- Silveri, M. C., Leggio, M. G., & Molinari, M. (1994). The cerebellum contributes to linguistic production: A case of agrammatic speech following a right cerebellar lesion. *Neurology, 44*, 2047–2050.
- Snyder, H. R., Banich, M. T., & Munakata, Y. (2011). Choosing our words: Retrieval and selection processes recruit shared neural substrates in left ventrolateral prefrontal cortex. *Journal of Cognitive Neuroscience, 23*, 3470–3482.

- Spinnler, H., & Tognoni, G. (1987). *Standardizzazione e taratura italiana di test neuropsicologici*. Milan: Masson Italia Periodici.
- Stoodley, C. J., & Schmahmann, J. D. (2009). The cerebellum and language: Evidence from patients with cerebellar degeneration. *Brain and Language, 110*, 149–153.
- Stoodley, C. J., Valera, E. M., & Schmahmann, J. D. (2010). An fMRI study of intra-individual functional topography in the human cerebellum. *Behavioural Neurology, 23*, 65–79.
- Stoodley, C. J., Valera, E. M., & Schmahmann, J. D. (2012). Functional topography of the cerebellum for motor and cognitive tasks: An fMRI study. *Neuroimage, 16*, 1560–1570.
- Strick, P. L., Dum, R. P., & Fiez, J. A. (2009). Cerebellum and nonmotor function. *Annual Review of Neuroscience, 32*, 413–434.
- Szaflarski, J. P., Allendorfer, J. B., Banks, C., Vannest, J., & Holland, S. K. (2013). Recovered vs. not-recovered from post-stroke aphasia: The contributions from the dominant and non-dominant hemispheres. *Restorative Neurology and Neuroscience, 31*, 347–360.
- Thompson-Schill, S. L., Swick, D., Farah, M. J., D'Esposito, M., Kan, I. P., & Knight, R. T. (1998). Verb generation in patients with focal frontal lesions: A neuropsychological test of neuroimaging findings. *Proceedings of the National Academy of Sciences, U.S.A., 95*, 15855–15860.
- Turkeltaub, P. E., Swears, M. K., D'Mello, A. M., & Stoodley, C. J. (2016). Cerebellar tDCS as a novel treatment for aphasia? Evidence from behavioral and resting-state functional connectivity data in healthy adults. *Restorative Neurology and Neuroscience, 34*, 491–505.
- Van Dun, K., Bodranghien, F. C. A. A., Mariën, P., & Manto, M. U. (2016). tDCS of the cerebellum: Where do we stand in 2016? Technical issues and critical review of the literature. *Frontiers in Human Neuroscience, 10*, 199.
- Zettin, M., Cappa, S., D'amico, A., Rago, R., & Perino, C. (1997). Agrammatic speech production after a right cerebellar haemorrhage. *Neurocase, 3*, 375–380.
- Zimmermann, P., & Fimm, B. (1994). *Tests d'évaluation de l'attention (TEA)*. Würselen, Germany: Psytest.